

+ THESE BANDS ARE AWESOME +
+ THESE BANDS ARE EVIL +

*THE YEAR OF 2006 / VIENNA



SUNN O)))

28.02.06, SZENE WIEN

the 1990s, the number of people who have been affected by the disease has increased significantly. In 1990, there were only 100 cases reported in the United States, but by 2000, the number had risen to over 10,000. This increase is due to a number of factors, including the fact that the disease is now being found in many more countries, and that it is becoming more common in people who are not in contact with the disease's natural host, the bat.

The disease is caused by a virus that is found in the saliva of bats. It is spread to humans when they are bitten by a bat, or when they come into contact with the bat's saliva. The disease is most common in people who live in areas where bats are common, such as caves and barns.

The disease is usually fatal, but there are some cases where people have recovered. However, the disease is still a major public health concern, and it is important to take steps to prevent its spread. This includes avoiding contact with bats, and using protective clothing when working in areas where bats are common.

The disease is also a concern for people who are immunocompromised, as they are more likely to develop the disease if they are exposed to the virus. It is important for these people to take extra precautions to avoid contact with bats, and to seek medical attention if they are bitten or come into contact with the bat's saliva.

The disease is also a concern for people who are traveling to areas where bats are common. It is important for these people to take precautions to avoid contact with bats, and to seek medical attention if they are bitten or come into contact with the bat's saliva.

The disease is also a concern for people who are working in areas where bats are common. It is important for these people to take precautions to avoid contact with bats, and to seek medical attention if they are bitten or come into contact with the bat's saliva.

The disease is also a concern for people who are living in areas where bats are common. It is important for these people to take precautions to avoid contact with bats, and to seek medical attention if they are bitten or come into contact with the bat's saliva.

The disease is also a concern for people who are visiting areas where bats are common. It is important for these people to take precautions to avoid contact with bats, and to seek medical attention if they are bitten or come into contact with the bat's saliva.

The disease is also a concern for people who are working in areas where bats are common. It is important for these people to take precautions to avoid contact with bats, and to seek medical attention if they are bitten or come into contact with the bat's saliva.



the 1990s, the number of people with a diagnosis of schizophrenia has increased in many countries, including the Netherlands (Kleinman and Lehmann 1995, van Os and van Dorp 1996). The prevalence of schizophrenia is estimated to be 1% in the general population (Kleinman and Lehmann 1995).

There is a growing awareness of the need to improve the quality of life of people with schizophrenia. The World Health Organization (WHO) has developed the concept of 'recovery' as a goal for people with schizophrenia (Kleinman and Lehmann 1995). Recovery is defined as a process of personal growth and development, leading to a more meaningful and satisfying life (Lehmann and Kleinman 1995).

One of the main goals of recovery is to help people with schizophrenia to live independently and to participate in society. This requires a range of services and supports, including housing, employment, and social activities. The Dutch government has developed a policy of 'deinstitutionalization', which aims to reduce the number of people in psychiatric hospitals and to increase the number of people living in the community (Kleinman and Lehmann 1995).

One of the main challenges of deinstitutionalization is to ensure that people with schizophrenia have access to the services and supports they need. This requires a range of services and supports, including housing, employment, and social activities. The Dutch government has developed a policy of 'deinstitutionalization', which aims to reduce the number of people in psychiatric hospitals and to increase the number of people living in the community (Kleinman and Lehmann 1995).

One of the main challenges of deinstitutionalization is to ensure that people with schizophrenia have access to the services and supports they need. This requires a range of services and supports, including housing, employment, and social activities. The Dutch government has developed a policy of 'deinstitutionalization', which aims to reduce the number of people in psychiatric hospitals and to increase the number of people living in the community (Kleinman and Lehmann 1995).

One of the main challenges of deinstitutionalization is to ensure that people with schizophrenia have access to the services and supports they need. This requires a range of services and supports, including housing, employment, and social activities. The Dutch government has developed a policy of 'deinstitutionalization', which aims to reduce the number of people in psychiatric hospitals and to increase the number of people living in the community (Kleinman and Lehmann 1995).

One of the main challenges of deinstitutionalization is to ensure that people with schizophrenia have access to the services and supports they need. This requires a range of services and supports, including housing, employment, and social activities. The Dutch government has developed a policy of 'deinstitutionalization', which aims to reduce the number of people in psychiatric hospitals and to increase the number of people living in the community (Kleinman and Lehmann 1995).

One of the main challenges of deinstitutionalization is to ensure that people with schizophrenia have access to the services and supports they need. This requires a range of services and supports, including housing, employment, and social activities. The Dutch government has developed a policy of 'deinstitutionalization', which aims to reduce the number of people in psychiatric hospitals and to increase the number of people living in the community (Kleinman and Lehmann 1995).





THE WILDERNESS

14.03.06, CHELSEA / WIEN





THE ENABLERS

27.04.06, CHELSEA / WIEN









TODD

27.04.06, CHELSEA / WIEN













